hydrobiologia volume 333 (1996), contents

volume 333, no. 1, october 7, 1996

Distribution of macroalgae and macroalgal communities in gradients of physical conditions in Potter Cove, King George Island, Antarctica
by H. Klöser, M.L. Quartino & C. Wiencke 1–17

The phytoplankton of some gravel-pit lakes in Spain
by M. Arauzo, M. Alvarez Cobelas, J. Vicioso & M. Verdugo 19–27

Reinstatement of the genus Paralona Sramek-Husek et al., 1962, and assessment of morphological variability in the Paralona pigra-group (Anomopoda: Chydoridae)
by H.J. Dumont & N.N. Smirnov 29–36

Patterns of nutrient release from nutrient diffusing substrates in flowing water
by L.D. Corkum 37–43

Planktonic cyclopoid copepods from small ponds in Kyushu, Japan. I. Subfamily Eucyclopinae with descriptions of micro-characters on appendages
by H. Ueda, T. Ishida & J-i. Imai 45–56

Effect of microbes on the uptake of ^{60}Co, ^{85}Sr, ^{95M}Tc, ^{131}I and ^{134}Cs by decomposing elm leaves in aquatic microcosms
by G.A. Bird & W. Schwartz 57–62

Analytical study of the locomotor and respiratory movements of tubificid worms by means of video recording
by C. Guérin & N. Giani 63–69

volume 333, no. 2, october 9, 1996

Food and feeding behaviour of a planktivorous cyprinid, *Pelecus cultratus* (L.), in a shallow eutrophic lake, Neusiedler See (Austria)
by Shengwen Liu & A. Herzig 71–77

Decomposition of *Eucalyptus globulus* leaves and three native leaf species (*Alnus glutinosa, Castanea sativa* and *Quercus faginea*) in a Portuguese low order stream
by C. Canhoto & M.A.S. Graça 79–85

Physiological reactions to a change in light regime in cultured *Skeletonema costatum* (Bacillariophyta): implications for estimation of phytoplankton biomass
by C. Brunet, D. Davoult & R. Casotti 87–94

The relative importance of water quality, sediment composition and floating vegetation in explaining the macrobenthic community structure of floodplain lakes (Paraná River, Argentina)
by J.A. Bechara 95–109

The epiphytic algae on floating macrophytes of a Paraná River floodplain lake
by G. Tesolín & G. Tell 111–120

The accumulation of amino-acids and humic substances in media conditioned by axenic and non-axenic duckweed (*Lemna minor* L.) and their possible ecological significance
by J.D. Thomas & P. Eaton 121–128

Chemistry, ecology and seasonal succession of Charophytes in the Al-Kharj Irrigation Canal, Saudi Arabia
by M.I. Hussain, T.M. Khoja & M. Guerlesquin 129–137

volume 333, no. 3, october 11, 1996

Distribution of stream macroalgae in the eastern Atlantic Rainforest of São Paulo State, southeastern Brazil
by C.C.Z. Branco & O. Necchi, Jr. 139–150
The relationship of floods, drying, flow and light to primary production and producer biomass in a prairie stream
by W.K. Dodds, R.E. Hutson, A.C. Eichem, M.A. Evans, D.A. Gudder, K.M. Fritz & L. Gray 151–159
Stylodrilus californianus n.sp., a new lumbriculid (Annelida: Oligochaeta) from North America
by P. Rodriguez 161–164
The effect of epibenthic predators and macroalgal cover on the benthic macroinvertebrate community of a shallow lagoon in the Bay of Cádiz (SW Spain)
by P. Drake & A.M. Arias 165–180
Microphytobenthos production in the Gulf of Fos, French Mediterranean coast
by C. Barranguet, M.R. Plante-Cuny & E. Alivon 181–193
Daphnia obtusa Kurz, 1874 emend Scourfield, 1942 from Australia
by J.A.H. Benzie & A.M.A. Hodges 195–199
Sulfate reduction in profundal sediments in Lake Kizaki, Japan
by J-h. Li, S. Takii, R. Kotakemori & H. Hayashi 201–208

Author index 209